

Mushroom immunomodulators: unique molecules with unlimited applications

Hesham A. El Enshasy^{1,2} and Rajni Hatti-Kaul³

¹ Institute of Bioproduct Development (IBD), Universiti Teknologi Malaysia (UTM), 81130 UTM, Skudai, Malaysia

² City of Scientific Research and Technology Application, New Burg Al Arab, Alexandria, Egypt

³ Department of Biotechnology, Center for Chemistry and Chemical Engineering, Lund University, Box 124, SE-221 00 Lund, Sweden

For centuries, mushrooms have been used as food and medicine in different cultures. More recently, many bioactive compounds have been isolated from different types of mushrooms. Among these, immunomodulators have gained much interest based on the increasing growth of the immunotherapy sector. Mushroom immunomodulators are classified under four categories based on their chemical nature as: lectins, terpenoids, proteins, and polysaccharides. These compounds are produced naturally in mushrooms cultivated in greenhouses. For effective industrial production, cultivation is carried out in submerged culture to increase the bioactive compound yield, decrease the production time, and reduce the cost of downstream processing. This review provides a comprehensive overview on mushroom immunomodulators in terms of chemistry, industrial production, and applications in medical and nonmedical sectors.

Introduction

Immunomodulators are key components in the modern health and wellness industries, reflecting the fact that the immune system is the first barrier for disease prevention. In any healthy organism, the immune system produces a wide range of immunomodulators to maintain homeostasis within the body. In clinical practice, immunomodulators are usually classified into immunosuppressants, immunostimulants, and immunoadjuvants (see [Glossary](#)). Their market share has increased rapidly over the past few years due to wide-ranging medical applications for stimulation and suppression of the immune system. They are even used as prodrugs or prophylactic medicine for healthy people. The market size of immunomodulators is valued at \$145.9 billion in 2012 and is proposed to increase rapidly with a compound

annual growth rate of 8.6% reaching \$259.3 billion in 2017 [1]. Chemically synthesized compounds and monoclonal antibodies of antiproliferative and antimetabolic drugs generate the highest revenues among all classes of immunomodulators. Recently, there has been growing interest in natural immunomodulators as alternatives to the currently used chemical drugs that have a high-risk profile [2]. Mushrooms are among the most interesting natural sources of compounds for pharmaceutical applications and are central components of traditional medicine worldwide. This is because they show greater diversity and possess unique bioactivity compared with other natural sources.

Mushroom polysaccharides such as lentinan, schizophyllan, polysaccharide K (Kerstin), and polysaccharide peptide (PSP) are now available on the pharmaceutical market. For example, lentinan is applied as adjuvant cancer immunotherapy or in parallel to radio- and chemothermotherapy [3].

Mushrooms: nutritional and medicinal facts

Mushrooms can be defined as macrofungi with distinctive fruiting bodies that are either epigeous (of fruiting bodies above the ground) or hypogeous (of underground fruiting bodies) and sufficiently conspicuous to the naked eye to be

Glossary

Fruiting body: is a multicellular structure on which spores producing structure as basidia or asci are born. They are distinct in size, shape, and coloration for each mushroom species.

Immunoadjuvants: are used to enhance vaccine efficacy. They can also be described as specific immune stimulators.

Immunomodulators: also known as biological response modifiers (BRMs), immunoaugmentors, or immunorestoratives, are substances, of biological or synthetic origin, which can stimulate, suppress or modulate any of the components of the immune system including both innate and adaptive arms of the immune response.

Immunostimulants: are agents that stimulate the immune system through induction or activation of immune system components or mediators. They enhance resistance against infection, allergy, cancer, and autoimmunity.

Immunosuppressants: are agents that inhibit the immune system. They can be used to control a pathological immune response after organ transplantation and to treat autoimmune diseases, hypersensitive immune reaction, and immunopathology associated with infections.

Immunotherapy: treatment (e.g., using drugs) that induces, enhances, or suppresses an immune response.

Mycelium: is the vegetative part of a fungus, consisting of a mass of branched hyphae. It forms the mushroom stalk in case of solid substrate grown mushrooms, or a complex shape of intertwined thread in submerged culture.

Nutraceuticals: food or part of food that provides medical or health benefits, including the prevention and treatment of disease.

Corresponding author: El Enshasy, H.A. (henshasy@ibd.utm.my).

Keywords: mushroom immunomodulators; lectins; fungal immunomodulatory proteins; polysaccharides; terpenes and terpenoids; medical application; animal feed and aquaculture; industrial production.

0167-7799/\$ – see front matter

© 2013 Elsevier Ltd. All rights reserved. <http://dx.doi.org/10.1016/j.tibtech.2013.09.003>



picked by hand [4]. Approximately 15 000 mushroom species have been identified worldwide, of which only about 650 species are reported to be of medicinal value [5]. Beside their high nutritional value, mushrooms are rich in many bioactive metabolites of high medicinal values such as lectins, polysaccharides, phenolic and polyphenolics, terpenoids, ergosterols, and volatile organic compounds [6]. Mushroom extracts have been shown to exhibit immunomodulator, antitumor/anticancer, antibacterial and antiviral, antioxidant, and antihypoglycemic activities, and as active medicine in the prevention of cardiovascular diseases by their effects as antiatherosclerosis agents [7,8]. Most medicinal mushroom research has been based on tests with crude extracts of the whole mushroom fruiting bodies or mycelia, or with partially purified bioactive compound.

Mushroom immunomodulators

More than 50 known mushrooms harbor immune-regulating organic compounds of highly diversified molecular weight and structure. Box 1 lists the top 10 mushrooms with immunomodulator activities. Mushrooms immunomodulators exhibit stimulating activities for both innate and adaptive immune systems. They proliferate and activate innate immune system components such as natural killer (NK) cells, neutrophils, and macrophages, and stimulate cytokines expression and secretion. These cytokines in turn activate adaptive immunity through the promotion of B cells for antibodies production and stimulation of T cell differentiation to T helper (Th) 1 and Th2 cells, which mediate cell and humoral immunities, respectively [9]. Based on their high molecular weight, mushroom polysaccharides are not able to penetrate the immune cells to activate immune cells directly. Thus, the stimulation mechanism of polysaccharides involves different cell receptors such as dectin-1, Complement receptor 3 (CR3), Lactosylceramide (LacCer), and Toll-like receptor (TLR)2. In such cases, the effectiveness of polysaccharides is governed by their binding affinity to immune cell receptors [10].

In general, mushroom immunomodulators are classified into four main groups: (i) immunomodulatory lectins; (ii) immunomodulatory terpenes and terpenoids; (iii) fungal immunomodulatory proteins (FIPs); and (iv) immunomodulatory polysaccharides. Each class is described below.

Lectins

Lectins form a diverse group of carbohydrate-binding proteins with specific binding capacities. They have been isolated from different organisms. However, those derived from mushrooms are characterized by specific immunomodulatory, antiproliferative, and antitumor activities. Lectins isolated from *Volvariella volacea* (straw mushroom) exhibit stronger immunomodulatory activity than other known lectins such as concanavalin A [11]. Two lectins extracted from *Tricholoma mongolicum*, TML-1 and TML-2, show immunomodulatory and antitumor activities when tested *in vivo* but not in *in vitro*, and they mediate their action through activation of the immune system rather than any direct cytotoxic effects [12]. These lectins stimulate the production of nitrite and tumor necrosis factor (TNF)- α , and inhibit the growth of mouse

Box 1. Top 10 mushrooms with immunomodulatory activity

1. *Agaricus subrufescens* (syn. *A. blazei* or *A. brasiliensis*). This edible mushroom has different common names [Almond portobello, Royal sun agaricus, Princess matsutake (JP)]. It is characterized by almond fragrance and a sweet taste. It contains immunostimulatory compounds such as β -1,3-D-glucans, glucomannan, and proteoglycans.
2. *Cordyceps sinensis* (syn. *Ophiocordyceps sinensis* or *Cephalosporium sinensis*) has been known for a thousand years as a medicinal mushroom in China and Taiwan. These pathogenic fungi are endoparasitic in arthropods. The immunomodulatory activity of this mushroom is due to the presence of polysaccharides and other compound such as cordycepin.
3. *Ganoderma lucidum*: Common names: Reishi (JP), Ligzhi (CN), spirit plant. It has a long history in Japanese and Chinese traditional medicine. This mushroom is rich with more than 50 types of polysaccharides and peptide-polysaccharides complexes in addition to about 120 bioactive compounds (mainly triterpenes). Most of these compounds can act as strong immunostimulants.
4. *Grifola frondosa*: Common name: Maitake. This mushroom is rich in β -glucan, which activates macrophages, NK cells and lymphokines. It also includes compounds such as ergosterol peroxide (EPO) and lipopolysaccharide (LPS) with significant immunomodulating activities.
5. *Hericium erinaceus*: Common names: Lion's Mane Mushroom, Bearded Tooth Mushroom, and Monkey Head Mushroom. It acts as an immunostimulant based on its content of bioactive polysaccharides of β -1,3-branched- β -1,2-mannan structure.
6. *Inonotus obliquus*: Common name: Chaga. This mushroom is a plant pathogen that grows on the trunk of birch trees in cold regions such as North America, Siberia, and Korea. The immunomodulating activity of this mushroom is based on its high content of β -glucans.
7. *Lentinula edodes*: Common name: Shiitake. This is an edible Japanese mushroom, famous for its immunomodulating properties. It is rich in a special type of polysaccharide called lentinan (1-3- β -D-glucan with 1-6- β -D-glucopyranoside branches).
8. *Pleurotus ostreatus*: Common name: Oyster mushroom. It is widely cultivated in different parts of the world. This mushroom includes different types of polysaccharides that belong to the β -glucan, heteroglycan, and proteoglycan families. Pleuran β -(1-3/1-6)-D-glucan is the most effective and well studied polysaccharide in this mushroom.
9. *Poria cocos*: Common names: Fu Ling (CN), Tuckahoe (US). This mushroom, used in traditional medicine, possesses immune stimulatory activity due to a special type of FIP, and *Poria cocos* protein (PCP).
10. *Trametes versicolor*: Common name: Turkey Tail. It has strong immunomodulatory effects due to a polysaccharide called Krestin (PSK), and also contains an FIP named tvC.

lymphoblast-like (p815) mastocytoma cells by the production of macrophage-activating factors. These factors include interferon (IFN)- γ and other cytokines, activated through upregulation of inducible nitric oxide synthase (NOS), interleukin (IL)-1 β , and transforming growth factor- β [13].

Lectins derived from the fruiting body of *Grifola frondosa* are characterized by their potent cytotoxic effects, even at very low concentrations, in HeLa cells. An 18-kDa thermostable lectin was isolated from *Ganoderma capense* with higher potent mitogenic activity than concanavalin A toward mouse splenocytes. It also showed antiproliferative activities on leukemia and hepatoma cells [14]. Recently, a novel 15.9-kDa homodimeric, lactose-binding, ricin-B-like lectin (CNL) was purified from the basidiomycete *Clitocybe nebularis* and exhibited antiproliferative activity against

human leukemic T cells [15]. This lectin induces the maturation and activation of dendritic cells (DCs) and thus stimulates several proinflammatory cytokines such as IL-6, IL-8, and TNF- α [16]. The CNL encoding gene has been cloned and successfully expressed in *Escherichia coli* [17].

Terpenes and terpenoids

Terpenes are a large and diversified group of organic compounds consisting of isoprene five-carbon atom units of molecular formula $(C_5H_8)_n$ as building blocks. The terpene compounds are named based on the number of repeated units of terpene building blocks, for example, monoterpenes (10 carbons), sesquiterpenes (15 carbons), diterpenes (20 carbons), sesterterpenes (25 carbons), triterpenes (30 carbons), and tetraterpenes (40 carbons). These compounds exist widely in plants as the main components of resin and essential oil. In macrofungi, terpenes are present in modified form (terpenoids or isoprenoids) and show biological activities with potential medical applications. Mushrooms belonging to *Ganoderma* sp., such as *Ganoderma lucidum* and *Ganoderma applanatum*, are known for their high content of triterpenoids, for example, lanostane, which shows immunomodulating and anti-infective activities [18]. The solvent extract of the fruiting body of this basidiomycete contains a wide range of terpenes and terpene derivatives such as ganodermic and ganoderic acids, ganoderals, ganoderols, ganodermantriol, lucidone, and ganodermanondiol (Figure 1), which exhibit biological activities including stimulation of the nuclear factor (NF)- κ B pathway and mitogen-activated protein kinases [19]. Recent research has shown that some *G. lucidum* terpenes can prevent drug nephrotoxicity and inflammation, suggesting that they may have pharmacological applications [20].

FIPs

In recent years, mushrooms have been reported to produce a new family of protein immunomodulators, termed FIPs. Since the discovery of the first FIP (Ling-Zhi-8 from *G. lucidum*) in 1989 [21], 11 different types of FIPs have

been isolated. These include FIP-fve from *Flammulina velutipes* (Gr.) Sing [22]; Ling-Zhi-8 (FIP-LZ-8 or FIP-glu) from *G. lucidum* [23]; FIP-gts from *Ganoderma tsugae* Murr [24]; FIP-gsi from *Ganoderma sinensis* [25]; FIP-pcp from *Poria cocos* (Schw.) Wolf [26]; and FIP-vvo and FIP-vvl from *Volvariella volvacea* (Bull.; Fr.) Sing [27]; FIP-aca from *Antrodia camphorate* [28]; FIP-gja from *Ganoderma japonicum*; FIP-gmi from *Ganoderma microsporium* [29]; in addition to one new FIP named FIP-tvc from *Trametes versicolor* [30]. These proteins are grouped together in one family based on highly similar amino acid sequences. They exist as dimers in a dumbbell-shaped structure similar to that of the variable region of immunoglobulin heavy chains, but they exhibit diverse activities (Table 1). The best-known FIP-LZ-8 protein consists of 110 amino acid residues and acts as an immunosuppressive agent. It has also been shown to suppress autoimmune diabetic reactions in an animal model and to increase graft survival in transplanted allogenic mouse skin, with fewer nephrotoxic effects compared to other immunosuppressive agents such as cyclosporine A [31].

The low yield of FIPs extracted from their native mushroom producer strain is a major limitation. Therefore, techniques are being developed for the production of recombinant FIPs in other organisms such as the yeast *Pichia pastoris* and the bacterium *E. coli*. Expression of the LZ-8 gene of *G. lucidum* in *P. pastoris*, for example, leads to the production of a recombinant LZ-8 protein (rLZ-8), which, although lacking the carbohydrate moiety of the native protein, shows the same bioactivity for IL-2 induction as the native protein [32]. The gene encoding FIP-fve protein is also expressed successfully in *E. coli* and the recombinant product exhibits activity similar to the native protein, inducing IL-2 and IFN- γ production [33]. More recently, two shuffled gene libraries were created for the genes *FIP-glu*, *FIP-gsi*, *FIP-fve*, *FIP-vvo* and the mutants *FIP-SN* and *FIP-SJ*, which were successfully expressed in *E. coli*. The recombinant FIPs showed higher immunomodulatory activities compared to those originally produced in mushrooms [34]. The shuffled FIPs, like the native proteins, induced the expression of specific cytokines (IL-2, IL-4, IFN- γ , IL-2 receptor in mouse splenocytes) but not TNF- α .




Figure 1. Mushroom triterpenes with immunomodulating activity. The immunomodulatory properties of ganoderic acid and ganodermanontriol are mediated through the activation of natural killer cells and enhancement of interleukin-2 and interferon- γ expression.

Polysaccharides

Mushrooms are an important source of different types of polysaccharides with immunomodulating activities [35]. Most of these polysaccharides are homoglycans (polysaccharides that contain residues of only one type of monosaccharide molecules) or heteroglycans (polysaccharides that contain residues of two or more types of monosaccharide molecules), and are able to combine with other proteins to make peptidoglycan or polysaccharide-protein complexes. The first reported polysaccharide with potential immunomodulating and anticancer activity was lentinan, a β -1,3-D-glucan with β -1,6 branches (Figure 2). This polysaccharide with triple helical structure was first isolated from the fruiting body of *Lentinus edodes* in the late 1960s in Japan [36]. Since then there have been many research efforts to discover new polysaccharide compounds with immunomodulating activities from mushroom fruiting body extracts. Until the late 1980s, only two more

Table 1. Some FIPs and their immunomodulatory activities

| Name of FIP | Producer organism | Immunomodulatory effect | Refs |
|-------------|------------------------------|---|------|
| FIP-aca | <i>Antrodia camphorata</i> | Induction of mRNA expression of different cytokines (IL-1 β , IL-6, IL-12, TNF- α) and chemokines (CCL3, CCL4, CCL5, CCL10) | [28] |
| FIP-fve | <i>Flammulina velutipes</i> | Stimulation of mitogenesis in human peripheral lymphocytes, suppression of systemic anaphylaxis reaction, enhanced transcription of IL-3, INF- γ | [22] |
| FIP-glu | <i>Ganoderma lucidum</i> | Enhanced transcription of IL-2, IL-3, IL-4, IFN- γ , TNF- α | [23] |
| FIP-gmi | <i>Ganoderma microsporum</i> | Down regulation of TNF- α | [29] |
| FIP-gts | <i>Ganoderma tsugae</i> | Induction of cytokine secretion, cellular proliferation of human peripheral mononuclear cells (HPBMCs), enhanced IFN- γ expression | [24] |
| FIP-gsi | <i>Ganoderma sinensis</i> | Enhanced production of IL-2, IL-3, IL-4, INF- γ , TNF- α | [25] |
| FIP-pcp | <i>Poria cocos</i> | Enhanced production of IL-1 β , IL-6, IL-18, TNF- α , NO | [26] |
| FIP-tve | <i>Trametes versicolor</i> | Increased human peripheral blood lymphocytes, enhanced production of TNF- α , NO | [30] |
| FIP-vvo | <i>Volvariella volvacea</i> | Enhanced expression of IL-2, IL-4, IFN- γ , TNF- α | [27] |

Abbreviation: CCL, chemokine CC ligand.

polysaccharides of β -glucan type, schizophyllan from *Schizophyllum commune* and the protein-bound polysaccharide Krestin from *Coriolus versicolor*, were characterized with fully proven chemical structure and activities [35]. They were successfully introduced to the nutraceutical and pharmaceutical market as biological response modifier (BRM) drugs based on their ability to restore or enhance immune responses in both *in vitro* and *in vivo* models. However, our understanding of the immunomodulatory effects of mushroom extracts in terms of chemical structure, and exact effects on the immune system remains incomplete.

Table 2 provides a short list of immunomodulator polysaccharides and polysaccharide–proteins of mushroom origin. In general, mushroom polysaccharides are highly diversified in their sugar composition, main chain polymer structure, degree of branching, conformation, molecular weight, and other physical properties, which together have significant effects on the bioactivity and mode of action of the polysaccharide, as discussed below [37].

Effect of molecular weight

In general, high molecular weight polysaccharides usually exhibit higher bioactivity. For example, in a (1 \rightarrow 3)- β -glucan extract of *Gr. frondosa* containing different molecular weight fractions, the highest immunomodulatory activity was ascribed to the fraction of 800 kDa and above [38]. In another study, when the polysaccharide krestin (PSK) was fractionated by ultrafiltration, the highest immunomodulatory activity was associated with the highest molecular weight fraction of >200 kDa [39]. By contrast, a low molecular weight 5-kDa fraction of hydrolyzed scleroglucan (a branched β -glucan from *Sclerotium rolfsii*) exhibited strong immunostimulatory activity, increasing the secretion of TNF- α and stimulating the proliferation of lymphocytes [40]. Low molecular weight polysaccharides can penetrate immune cells and exert stimulatory effects. However, the superiority of high molecular weight polysaccharides (which are not able to penetrate the cells) was not clearly discussed in the literature. The superior activity of high molecular weight polysaccharides may be attributed to the better binding affinity to the carbohydrate receptors of the immune cells.

Effect of branching

In most cases, the bioactive immunomodulator polysaccharides are characterized by a main chain of 1-3,- β -D-glucan with a small number of short branched chains with 1-6, β -linkage. The most active polymers generally have a degree of branching number (DB) between 20% and 33%. For example, lentinan (DB = 40% or 2/5) and schizophyllan (DB = 33% or 1/3) are both 1-3,- β -D-glucans with one and two branches for every 5-D-glucopyranosyl and 3-D-glucopyranosyl residue, respectively. The polysaccharide moiety of PSK (DB = 20% or 1/5) is composed of (1-3)- β -(1-4)- β -D-glucan with one branch for every 5-D-glucopyranosyl residue. Even complete debranching of polysaccharides sometimes increases their bioactivity, for example, pachyman, a branched 1-3,- β -D-glucan of the brown rot fungus *Poria cocos* is inactive, whereas the debranched form, pachymanan, obtained after mild selective periodate oxidation, shows significant activity [41].

Debranching of lentinan also increases its biological activity as immunostimulant and antitumor agent [42]. More recently, a study of the correlation between branching and bioactivity of lentinan showed that among different polysaccharides fractions with DB ranging between 19 and 50%, the maximal immunomodulating and antitumor activities were achieved when the molecule had a DB of 32% [43]. Therefore, we can conclude that the relation between branching and polysaccharides immunomodulatory activity is case specific and needs further investigation.

Effect of helical conformation

Mushroom polysaccharides can exist as single or triple helical forms, as well as random coiled structures. The triple helix conformation is usually more stable than the other structures and bears the cytokine stimulating activity of the β -D-glucan. The most bioactive mushroom polysaccharides known, lentinan, schizophyllan, scleroglucan, and the glucan moiety of PSK, all have a triple helix structure [44]. Furthermore, among different polysaccharide fractions isolated from *Herichium erinaceus*, a low molecular weight fraction with triple helix conformation of a β -1,3-branched- β -1,6-glucan exhibited immunostimulating activity that included macrophage activation, NO production, and expression of IL-1 β and TNF- α [45]. On the contrary, other studies have shown that mushroom




Figure 2. Some examples of polysaccharides produced by mushroom of immunomodulating activities. **(A)** Pleuran (CAS No. 159940-37-1) is an insoluble polysaccharide [β -(1,3/1,6)-D-glucan], isolated from *Pleurotus ostreatus*. **(B)** Lentinan (CAS No. 37339-90-5) is a polysaccharide isolated from the fruiting body of shiitake (*Lentinula edodes*). Lentinan is a β -1,3 glucan with β -1,6 branching.

polysaccharides with random coil conformation can also exhibit potent immunomodulating and anticancer activity [46].

Chemical modifications of polysaccharides

With an aim to increase the biological activity, chemical modification of the polysaccharides to produce carboxymethylated, hydroxylated, formylmethylated, aminoethylated, or sulfated derivatives has been attempted. The hydroxylated form of schizophyllan, for example, was reported to stimulate the production of NO and TNF- α in macrophages better than the native polysaccharides [47]. Also the sulfated (1-3)- α -D-glucan, a native cell wall glucan from *L. edodes*, exhibited higher immunomodulatory and anticancer activity compared to the native polysaccharides [48]. Another example is the improved immunostimulating activity after sulfonation of (TAPA1), a polysaccharide extracted from *Tremella aurantialba* fruiting bodies [49]. A carboxymethylated-sulfated β -(1-3)-D-glucan from *Po. cocos* showed five times greater anticancer activity compared to the native polysaccharides.

The induction of carboxymethyl and sulfate groups increased the possible contact between the modified polysaccharides with the immune cell receptor through hydrogen binding and electrostatic attraction and thus increased the immunological response. This was also proven by the increase in phagocyte and thymus indexes, spleen index, and antibody production. [50]. However, other researchers reported that the high stimulatory effect of the hyper branched β -glucan TM3b after sulfation was mainly attributed to the increase in solubility of the molecule [51]. Thus molecular weight, branching, chemical configuration, and chemical modification have strong influence on the solubility of the polysaccharide, which also reflect directly on molecule bioactivity. Thus, all these polysaccharide features should be studied in an inter-related context for better understanding of the immunomodulatory potency of the compounds.

Medical applications

In spite of many *in vitro* and *in vivo* studies performed during the past 50 years revealing the potential of

Table 2. Mushroom polysaccharides and their immunomodulator activities

| Mushroom Latin name | Common names | Biomaterial | Active compound | Immunomodulatory activity | Refs |
|---|---|-------------|---|--|------|
| <i>Agaricus blazei</i> (<i>Agaricus subrufescens</i>) | Royal sun agaricus, Almond portobello | FB, CB | Glycoprotein (ATOM), β -1, 3-D-glucan, with β -1,6-D-glucan branch | Induction of TNF, IFN- γ , and IL-8 production | [68] |
| <i>Antrodia camphorata</i> | Poroid brown-rot fungus | M | β -1,3-D-Gluco-pyranans with β -1,6-D-glucosyl branches, proteoglycan | Induction of INF- γ , TNF- α | [69] |
| <i>Cordyceps sinensis</i> | Caterpillar fungus | FB, SM, CB | β -D-glucan, heteroglycan, cordyglucan | Increase in IL-5 induction with decrease in IL-4 and IL-17 | [70] |
| <i>Cryptoporus volvatus</i> | Grey-Brown Sap Rot | FB | β -1,3-D-Glucan | Decrease in TLR2 and activate NF- κ B | [71] |
| <i>Flammulina velutipes</i> | Golden needle mushroom | FB, SM, CB | Glycoprotein, peptidoglycan, (FVP) | Increase NO, IL-1 production, and TNF- α secretion | [72] |
| <i>Ganoderma lucidum</i> | Reishi, Ling-zhi, Spirit plant | FB, CB | Ganoderan, Heteroglycan, mannoglucan, glycopeptide | Stimulate TNF- α , IL-1, IFN- γ production, activate NF- κ B. | [73] |
| <i>Grifola frondosa</i> | Hen of the woods, Maitake | FB, CB, SM | Grifolan (1-6-monoglucosyl- branched β -1,3-D-glucan), proteoglycan, heteroglycan, galactomannan | Macrophage activation, induction of IL-1, IL-6, and TNF- α secretion | [74] |
| <i>Hericium erinaceus</i> | Lion's Mane Mushroom, Bearded Tooth Mushroom, Monkey head Mushroom | FB, CB | Heteroglycan, heteroglycan- peptide, β -1,3 branched- β -1,2-mannan | Induce NO production, increase expression of TNF- α , IL-1 β , IL-12 | [45] |
| <i>Inonotus obliquus</i> | Clinker polypore, birch conk, Chaga | FB, SM | β -D-glucan | Enhance expression of IL-1 β , IL- 6, TNF- α , and iNOS in macrophages | [75] |
| <i>Lentinus edodes</i> | Shiitake, black forest mushroom, golden oak mushroom | FB, CB | Lentinan, glucan, mannoglucan, proteoglycan | Induces non-specific cytotoxicity in macrophage and enhance cytokine production | [76] |
| <i>Morchella esculenta</i> | Morel mushroom, sponge morel | FB | Galactomannan, β -1,3-D-glucan | Macrophage activation, activate NF- κ B | [77] |
| <i>Morchella conica</i> | Sponge mushroom | FB, CB | Galactomannan | Induces NO, IL-1 β , IL-6 production | [78] |
| <i>Phellinus linteus</i> | Mesima, Black Hoof Fungus | FB | Acidic polysaccharides | Activation of murine B cells, Induce IL-12 and IFN- γ production Block NF- κ B, TNF- α , IL-1 α , IL-1 β , and IL-4 production | [79] |
| <i>Pleurotus ostreatus</i> | Oyster mushroom | FB, SM, CB | Pleuran, heterogalactan, proteoglycan | Induce IL-4 and IFN- γ production | [63] |
| <i>Polyporus umbellatus</i> (<i>Grifola umbellata</i>) | Umbrella polypore | SM, FB | Polysaccharides | Enhances TNF- α , IL-1 β , and NO production | [80] |
| <i>Polystrictus versicolor</i> | Turkey tail | FB, SM, CB | Krestin, heteroglycan, glycopeptide, polysaccharide K (PSK), polysaccharide peptide (PSP) | Stimulate T cell activation, induce IFN- γ and IL-2 production, induce gene expression of cytokines (TNF- α , IL-1, IL-6, IL-8) | [81] |
| <i>Sarcodon aspratus</i> | Black tiger paw | FB | Fucogalactan, 1,6- α -D- glucopyranosyl residue | Increase the release of TNF- α and NO in macrophage | [82] |
| <i>Schizophyllum commune</i> | Split Gill | FB, WM | Schizophyllan, 1,6- monoglucosyl branched β -1, 3-D-glucan | Activation of T cell, increase interleukin, and TNF- α production | [83] |
| <i>Sclerotinia sclerotiorum</i> | White mold | FB | Scleroglucan (1,6- monoglucosyl-branched β -1, 3-D-glucan) | Stimulates lymphocyte proliferation, increase release of TNF- α | [40] |
| <i>Sparassis crispa</i> | Rooting cauliflower mushroom | FB | β -Glucan | Enhances IL-6 and INF- γ production | [84] |
| <i>Tremella aurantialba</i> | The golden Tremella | FB | Heteroglycan | Enhances mouse spleen lymphocyte proliferation | [49] |
| <i>Xylaria nigripes</i> | Dead moll's fingers | WM | β -Glucan | Inhibits NO, IL-1 β , IL-6, TNF- α , and IFN- γ production | [85] |

Abbreviation: M, mycelium; WM, whole mushroom; SM, submerged mycelium; CB, culture broth; FB, fruiting bodies.

mushroom extracts as immunomodulators, little information is available on their efficacy in clinical trials. The mushroom polysaccharides of defined chemical structure such as lentinan, schizophyllan, and grifolan have

been used in clinical trials. Lentinan was the first approved mushroom polysaccharide for clinical use as an immune adjuvant to chemotherapy for stomach cancer treatment in Japan since 1985 [52]. Nowadays, mushroom

polysaccharides are available in the market as nutraceuticals under different brand names such as Ganodex, Immuna, Lentinex, Immunoglukan, Pure Red Reishi Capsule, LifeShield, Bene-X, and Zymucan. These products are widely used as adjuvants to cancer radio- and chemotherapy, antibiotics, and vaccines to minimize their side

effects. Recent research shows that pleuran-based nutraceutical product (Immunoglukan P4H) reduces morbidity caused by recurrent respiratory tract infections through the modulation of humoral and cellular immunity [53].

The anticancer activity of mushroom polysaccharides is mediated through the activation of the immune system




Figure 3. Process for pleuran production by *Pleurotus ostreatus* using submerged mycelium fermentation (Process A) or solid state fermentation (Process B). Process A begins with mushroom cell propagation on agar plates followed by submerged cultivation of cells in shake flasks, as an intermediate step for inoculum preparation for bioreactor culture. Cultivation in the bioreactor is carried out by scaling up the process from a small to a large bioreactor followed by centrifugation to separate cells from the culture broth. If the polysaccharides produced are extracellular, extraction is usually carried out either through direct precipitation using cold ethanol followed by lyophilization to obtain the crude extract. A solvent-free method was also developed to separate high molecular weight compounds directly by ultrafiltration followed by direct lyophilization of the polysaccharides. The endopolysaccharides are extracted using the conventional scheme of polysaccharides extraction from mushroom fruiting bodies (Process B). With fruiting bodies as starting materials (Process B), cells are first washed and dried followed by homogenization prior to several hours of extraction with boiling water. The water extract of crude polysaccharide is concentrated using vacuum extraction followed by alcohol precipitation using cold ethanol. The precipitate is then lyophilized. In this process, further purification could be also done by repeating the steps between water extraction step and lyophilization.

[46,54]; however, other mechanisms such as direct inhibition of cancer cell growth (apoptotic and necrotic effect) can also contribute. In addition to the above applications, mushroom immunomodulators are currently used in cosmeceutical industries and have become basic ingredients in functional creams for wound repair and antiaging [55].

Applications in animal feed and aquaculture

Different types of antibiotics have been used widely as integral parts of animal feed systems to improve growth, control microbial infections, and decrease mortality during the spread of epidemic disease and the early stage of animal growth. However, with the increasing threat of antibiotic-resistant pathogens, their application has been banned in Europe since 2006 [Regulation (EC) No. 1831/2003] [56]. Therefore, replacement of antibiotics with natural immunomodulators to boost natural defense has been of interest [57]. Being a high volume–low value market, most research has focused on using whole mushroom or crude extracts as animal or aquaculture feed additives. Recently, mushrooms were added as feed additives either alone or in combination with probiotics to enhance the immune system in broiler chickens [58].

The immunostimulatory effect of mushroom was mainly attributed to the enhancement of probiotic flora growth, limiting the growth of microbial pathogens. Shiitake mushroom extract has been used to boost the immune system of chickens through direct activation of lymphocytes and macrophages; the level of mRNAs encoding IL-1 β , IL-6, IL-12, and IL-18 in macrophages was enhanced [59]. In aquaculture application, diet enrichment with the black hoof fungus (*Phellinus linteus*) extract enhanced the disease resistance of kelp grouper fish against pathogens such as *Vibrio harveyi*, *Vibrio alginolyticus*, and *Vibrio carchariae*; the resistance was mediated through enhancement of cellular and humoral immune response components such as serum lysozyme activity, and phagocytic activity [60]. The immunostimulatory potential of lentinan extract for healthy growth of sea cucumber (*Apostichopus japonicus*) was reported recently, which was mediated through the increase of phagocytic activity and viability of coelomocytes [61]. A significant increase in immune response and *Vibrio* disease resistance of white shrimp (*Litopenaeus vannamei*) was obtained using feed supplemented with small particle of monkey head mushroom *Hericium erinaceum*. This was due to the increase in phenoloxidase, superoxide dismutase and glutathione peroxidase activities. Recently, it was shown that administration of β -glucan mixture enhances the innate immune response in the high value orange spotted grouper fish (*Epinephelus coioides*) and protects it against *V. alginolyticus* [62].

Industrial production of mushroom immunomodulators

Mushrooms were first cultivated using hardwood tree logs as a substrate; producing high levels of lignocellulosic hydrolytic enzymes, mushrooms are able to use wood components as nutrition source. This process occurs over several years and yields only two crops per year. Subsequently, solid state fermentation (SSF), in which mushrooms grow on a combination of compost, wood residue, and minerals in polypropylene bags, became more common in

mushroom farms. After autoclaving, the bags are inoculated by actively growing mycelia and incubated in greenhouses. After a long cultivation time, sometimes exceeding 2 months, the fruiting bodies are harvested and used for extraction and purification of the bioactive compound. In SSF, the substrates can vary based on mushroom type. In this process, temperature and humidity should be carefully controlled, and the greenhouse is kept clean to reduce the risk of contamination with pathogens. However, for production of mushroom immunomodulators for medicinal applications, the process must be in compliance with current good manufacturing practice (cGMP) for active pharmaceutical ingredients (APIs), which is difficult to implement in SSF given that it is an open cultivation system; there are many challenges including the quality of substrate materials, cultivation parameters and controlled yield of the desired compound. Thus, in recent years research has focused on mushroom cultivation in a closed system using submerged fermentation (SMF). Its advantages include a higher yield of active metabolite in a shorter time, total control of all cultivation parameters, cultivation under fully sterile conditions, and easier downstream processes. Figure 3 shows a simple production platform for mushroom polysaccharides using SSF and SMF systems. Like other fungal fermentations, this process is governed by several factors including medium composition, cultivation conditions such as pH, temperature, aeration rate, and agitation, and fungal morphology in the cultivation vessel [63–66]. The type and concentration of carbon and nitrogen sources applied, C/N ratio, and addition of trace elements/supplements strongly affect polysaccharide biosynthesis. Moreover, oxygen supply acts as one of the bottleneck parameters affecting the yield of polysaccharides production. Fed-batch cultivation seems to be the most appropriate method for high polysaccharides yield as in case of *Agaricus brasiliensis* and *Pleurotus ostreatus* [67].

Concluding remarks

This review has highlighted the importance of mushrooms as future biofactories for the production of immunomodulator compounds of highly diverse chemical structures with potential applications in human and animal health. However, the application of mushroom immunomodulators in modern medicine faces six main challenges that need to be addressed. First, most mushrooms are difficult to cultivate in greenhouses; their availability is seasonal and weather dependent. Second, the content of bioactive ingredients depends on the collection time, procedure, season, and environment. Third, the mushroom greenhouse cultivation method is not validated and not run according to the cGMP requirements. Therefore, more research is needed for the development of cultivation processes in submerged culture to produce bioactive metabolites for pharmaceutical applications. Fourth, mushrooms are slow growing and produce bioactive compounds in low concentrations. This could be overcome by expressing the relevant genes in other hosts with higher growth rate and productivity such as *E. coli*, *P. pastoris*, and the fungus *Aspergillus niger*. Fifth, in most of the research conducted thus far, the immunomodulatory activities were assayed using crude mushroom extracts or

mixtures of different metabolites. It will be necessary to isolate and identify the active metabolites for better understanding of the immunomodulatory mechanism of each particular compound. Upgrading the crude extract from the nutraceutical to the pharmaceutical market will require careful product formulation and clinical trials to determine the proper dose and prove efficacy. Finally, there is a lack of standard testing protocols to guarantee the quality and the efficacy of the mushroom products for pharmaceutical applications. Some mushroom compounds, such as polysaccharides, are highly diversified in terms of structure and molecular weight; thus, it is difficult to produce the same quality from batch to batch. Increasing demand for these natural products will drive development towards overcoming these barriers for entering the market.

References

- 1 GBI Research (2012) *Immunomodulators Market to 2017–Immunosuppressant Antibodies to Replace Antimetabolites as the Highest Value Sector by 2017* (<http://www.marketresearch.com/GBI-Research-v3759/Immunomodulators-Immunosuppressant-Antibodies-Replace-Antimetabolites-6784727/>)
- 2 Shukla, S. *et al.* (2012) Plants as potential sources of natural immunomodulators. *Rev. Environ. Sci. Biotechnol.* <http://dx.doi.org/10.1007/s11157-012-9303-x>
- 3 Lindequist, U. *et al.* (2005) The pharmacological potential of mushrooms. *Evid. Based Complement. Altern. Med.* 2, 285–299
- 4 Chang, S.T. and Miles, P.G. (1992) Mushroom biology – a new discipline. *Mycologist* 6, 64–65
- 5 Minato, K. (2010) Mushrooms: immunomodulating activity and role in health promotion. In *Dietary Components and Immune Function* (Watson, R.R. *et al.*, eds), Springer Science+Business Medium
- 6 Kalač, P. (2013) A review of chemical composition and nutritional value of wild-growing and cultivated mushrooms. *J. Sci. Food Agric.* 93, 209–218
- 7 Zong, A. *et al.* (2012) Anticancer polysaccharides from natural sources: a review of recent research. *Carbohydr. Polym.* 90, 395–410
- 8 Mu, H. *et al.* (2012) Antioxidative properties of crude polysaccharides from *Inonotus obliquus*. *Int. J. Mol. Sci.* 13, 9194–9206
- 9 Borchers, A.T. *et al.* (2008) The immunobiology of mushrooms. *Exp. Biol. Med.* 233, 259–276
- 10 Chen, J. and Seviour, R. (2007) Medicinal importance of fungal β -(1→3), (1→6)-glucan. *Mycol. Res.* 111, 635–652
- 11 Sze, S.C. *et al.* (2004) *Volvariella volvacea* lectin activates mouse T lymphocytes by a calcium dependent pathway. *J. Cell. Biochem.* 92, 1193–1202
- 12 Wang, H.X. *et al.* (1997) Actions of lectins from the mushroom *Tricholoma mongolicum* on macrophages, splenocytes and life-span in sarcoma-bearing mice. *Anticancer Res.* 17, 419–424
- 13 Liu, F. *et al.* (2005) Lectin from *Tricholoma mongolicum* S. Imai (Agaricomycetidae) mycelia stimulates gene expression of immunomodulating cytokines in mouse peritoneal macrophages and splenocytes. *Int. J. Med. Mushrooms* 7, 243–248
- 14 Patrick, H. *et al.* (2004) A mushroom (*Ganoderma capense*) lectin with spectacular thermostability, potent mitogenic activity on splenocytes, and antiproliferative activity toward tumor cells. *Biochem. Biophys. Res. Commun.* 314, 988–993
- 15 Pohleven, J. *et al.* (2009) Purification, characterization and cloning of a ricin B-like lectin from mushroom *Clitocybe nebularis* with antiproliferative activity against human leukemic T cells. *Biochem. Biophys. Acta* 1790, 173–181
- 16 Švajger, U. *et al.* (2011) CNL, a ricin B-like lectin from mushroom *Clitocybe nebularis*, induces maturation and activation of dendritic cells via the toll-like receptor 4 pathway. *Immunology* 134, 409–418
- 17 Pohleven, J. *et al.* (2012) Bivalent carbohydrate binding is required for biological activity of *Clitocybe nebularis* lectin (CNL), the N,N'-diacetylactosediamine (GalNAc β 1-4GlcNAc, LacdiNAc)-specific lectin from basidiomycete *C. nebularis*. *J. Biol. Chem.* 287, 10602–10612
- 18 Jeong, Y.T. *et al.* (2008) *Ganoderma applanatum*: a promising mushroom for antitumor and immunomodulating activity. *Phytother. Res.* 22, 614–619
- 19 Gao, Y. *et al.* (2003) A phase I/II study of a *Ganoderma lucidum* (Curt:Fr.) P. Karst. Extract (ganopoly) in patients with advanced cancer. *Int. J. Med. Mushrooms* 4, 207–214
- 20 Pillai, T.G. *et al.* (2011) Prevention of cisplatin induced nephrotoxicity by terpenes isolated from *Ganoderma lucidum* occurring in southern parts of India. *Exp. Toxicol. Pathol.* 63, 157–160
- 21 Tanaka, S. *et al.* (1989) Complete amino acid sequence of an immunomodulatory protein, ling zhi-8 (LZ-8) an immunomodulator from a fungus *Ganoderma lucidum*, having similarity to immunoglobulin variable regions. *J. Biol. Chem.* 464, 16372–16377
- 22 Wang, P.-H. *et al.* (2004) Immunomodulatory protein from *Flammulina velutipes* induces interferon- γ production through p38 mitogen-activated protein kinase signaling pathway. *Agric. Food Chem.* 52, 2721–2725
- 23 Li, Q.Z. *et al.* (2010) *In vitro* synthesis of a recombinant fungal immunomodulatory protein from Lingzhi or Reishi medicinal mushroom, *Ganoderma lucidum* (W. Curt.: Fr.) P. Karst. (Aphyllophoromycetidae) and analysis of its immunomodulatory activity. *Int. J. Med. Mushrooms* 12, 347–358
- 24 Hsiao, Y.-M. *et al.* (2008) Effect of a fungal immunomodulatory protein from *Ganoderma tsugae* on cell cycle and interferon-gamma production through phosphatidylinositol 3-kinase signal pathway. *Process Biochem.* 43, 423–430
- 25 Li, Q. *et al.* (2010) Cytokines expression induced by *Ganoderma sinensis* fungal immunomodulatory proteins (FIP-gsi) in mouse spleen cells. *Appl. Biochem. Biotechnol.* 162, 1403–1413
- 26 Chang, H.-H. and Sheu, F. (2007) A novel fungal immunomodulatory protein (PCP) isolated from *Poria cocos* activates mouse peritoneal macrophage involved in toll-like receptor 4. *FASEB J.* 21, 702–715
- 27 Hsu, H.-Y. *et al.* (2008) Reishi immuno-modulation protein induces interleukin-2 expression via protein kinase-dependent signaling pathways within human T-cells. *J. Cell. Physiol.* 215, 15–26
- 28 Sheu, F. *et al.* (2009) Purification, cloning, and functional characterization of a novel immunomodulatory protein from *Antroidea camphorate* (Bitter mushroom) that exhibits TLR2-dependent NF- κ B activation and M1 polarization within murine macrophages. *J. Agric. Food Chem.* 57, 4130–4141
- 29 Lin, C.H. *et al.* (2010) GMI, a *Ganoderma* immunomodulatory protein, down-regulates tumor necrosis factor α -induced expression of matrix metalloproteinase 9 via NF- κ B pathway in human alveolar epithelial A549 cells. *J. Agric. Food Chem.* 58, 12014–12021
- 30 Fend, L.I. *et al.* (2011) Purification and characterization of a novel immunomodulatory protein from the medicinal mushroom *Trametes versicolor*. *Sci. China Life Sci.* 54, 379–385
- 31 Van der Hem, L.G. *et al.* (1996) Ling Zhi-8: a fungal protein with immunomodulatory effects. *Transplant. Proc.* 28, 958–959
- 32 Hsu, H.-C. *et al.* (1997) Fip-voo, a new fungal immunomodulatory protein isolated from *Volvariella volvacea*. *Biochem. J.* 323, 557–565
- 33 Kong, X. *et al.* (2013) High-yield production in *Escherichia coli* of fungal immunomodulatory protein isolated from *Flammulina velutipes* and its bioactivity assay *in vivo*. *Int. J. Mol. Sci.* 14, 2230–2241
- 34 Wang, X.-F. *et al.* (2013) *In vitro* rapid evolution of fungal immunomodulatory proteins by DNA family shuffling. *Appl. Microbiol. Biotechnol.* 97, 2455–2465
- 35 El Enshasy, H. (2010) Immunomodulators. In *The Mycota: Industrial Applications* (2nd edn) (Hoffrichter, M., ed.), pp. 165–194, Springer Verlag
- 36 Ikekawa, T. *et al.* (1969) Antitumor activity of aqueous extracts of edible mushrooms. *Cancer Res.* 29, 734–735
- 37 Ooi, V.E.C. (2008) Antitumor and immunomodulatory activities of mushroom polysaccharides. In *Mushrooms as Functional Food* (Cheung, P.C.K., ed.), pp. 147–198, Wiley
- 38 Adachi, Y. *et al.* (1990) Change of biological activities of (1→3)- β -D-glucan from *Grifola frondosa* upon molecular weight reduction by heat treatment. *Chem. Pharm. Bull.* 38, 477–481
- 39 Kim, F. *et al.* (1990) Stimulation of interferon- γ -induced human myelogenous leukemic cell differentiation by high molecular weight PSK subfraction. *Anticancer Res.* 10, 55–58
- 40 Bimczok, D. *et al.* (2009) Short chain regioselectively hydrolyzed scleroglucans induce maturation of porcine dendritic cells. *Appl. Microbiol. Biotechnol.* 82, 321–331

- 41 Chihara, G. *et al.* (1970) Antitumor polysaccharide derived chemically from natural glucan (pachyman). *Nature* 225, 943–944
- 42 Ren, L. *et al.* (2010) Antitumor activity of mushroom polysaccharides: a review. *Food Funct.* 3, 1118–1130
- 43 Bae, I.Y. *et al.* (2013) Correlation of branching structure of mushroom β -glucan with its physiological activities. *Food Res. Int.* 51, 195–200
- 44 Palleschi, A. *et al.* (2005) Molecular dynamics investigations of the polysaccharide scleroglucan: first study on the triple helix structure. *Carbohydr. Res.* 340, 2154–2162
- 45 Lee, J.S. *et al.* (2009) Study on macrophage activation and structural characteristics of purified polysaccharides from the liquid culture broth of *Hericium erinaceus*. *Carbohydr. Polym.* 78, 162–168
- 46 Wasser, S.P. (2002) Medical mushrooms as a source of antitumor and immunomodulating polysaccharides. *Appl. Microbiol. Biotechnol.* 60, 258–274
- 47 Ohno, N. *et al.* (1995) Comparison of the immunopharmacological activities of triple and single-helical schizophyllan in mice. *Biol. Pharmaceut. Bull.* 18, 1242–1247
- 48 Zhang, P. and Cheung, P.C.K. (2002) Evaluation of sulfated *Lentinus edodes* α -(1 \rightarrow 3)-D-glucan as a potential antitumor agent. *Biosci. Biotechnol. Biochem.* 66, 1052–1056
- 49 Du, X-J. *et al.* (2010) Purification, chemical modification and immunostimulating activity of polysaccharides from *Tremella aurantialba* fruit bodies. *J. Zhejiang Univ. Sci. B* 11, 437–442
- 50 Chen, X. *et al.* (2010) Immunopotential and anti-tumor activity of carboxymethylated-sulfated β -(1-3)-D-glucan from *Poria cocos*. *Int. Immunopharmacol.* 10, 398–405
- 51 Tao, Y. *et al.* (2006) Physicochemical properties and antitumor activities of water soluble native and sulfated hyperbranched mushroom polysaccharides. *Carbohydr. Res.* 341, 2261–2269
- 52 Higashi, D. *et al.* (2012) The effect of lentinan combination therapy for unresectable advanced gastric cancer. *Anticancer Res.* 32, 2365–2368
- 53 Jesenak, M. *et al.* (2013) Immunomodulatory effect of pleuran (β -glucan from *Pleurotus ostreatus*) in children with recurrent respiratory tract infections. *Int. Immunopharmacol.* 15, 395–399
- 54 Vannucci, L. *et al.* (2013) Immunomodulatory properties and antitumor activities of glucans (Review). *Int. J. Oncol.* 43, 357–364
- 55 Hyde, K.D. *et al.* (2010) Fungi—an unusual source for cosmetics. *Fungal Divers.* 43, 1–9
- 56 Regulation (EC) N. 1831/2003 of the European Parliament and of the Council of 22 september 2003 on Additives for Use in Animal Nutrition (<http://irmm.jrc.ec.europa.eu/SiteCollectionDocuments/EC-1831-2003.pdf>)
- 57 Gallois, M. *et al.* (2009) Natural alternatives to in-feed antibiotics in pig production: can immunomodulators play a role? *Animal* 3, 1644–1661
- 58 Willis, W.L. *et al.* (2009) Administration of mushroom extract to broiler chickens for bifidobacteria enhancement and *Salmonella* reduction. *J. Appl. Poult. Res.* 18, 658–664
- 59 Lee, S.H. *et al.* (2010) *In vitro* effects of plant and mushroom extracts on immunological function of chicken lymphocytes and macrophages. *Br. Poult. Sci.* 51, 213–221
- 60 Harikrishnan, R. *et al.* (2011) Diet enriched with mushroom *Phellinus linteus* extract enhances the growth, innate immune response, and disease resistance of kelp grouper, *Epinephelus bruneus* against vibriosis. *Fish Shellfish Immunol.* 30, 128–134
- 61 Wang, Y. *et al.* (2013) Lentinan extracted from shiitake mushrooms (*Lentinus edodes*) improves the non-specific immunity of sea cucumber (*Apostichopus japonicas*). *Aquacult. Int.* <http://dx.doi.org/10.1007/s10499-013-9628-7>
- 62 Chang, C-S. *et al.* (2013) Innate immune responses and efficacy of using mushroom beta-glucan mixture (MBG) on orange-spotted grouper, *Epinephelus coioides*, aquaculture. *Fish Shellfish Immunol.* 35, 115–125
- 63 El Enshasy, H.A. *et al.* (2012) Pleuran: immunomodulator polysaccharide from *Pleurotus ostreatus*, structure, production and application. In *Mushrooms: Types, Properties and Nutrition* (Andres, S. and Baumann, N., eds), pp. 153–172, Nova Publisher
- 64 Huang, H-C. and Liu, Y-C. (2008) Enhancement of polysaccharide production by optimization of culture conditions in shake flask submerged cultivation of *Grifola umbellata*. *J. Chin. Inst. Chem. Eng.* 39, 307–311
- 65 Papaspyridi, L-M. *et al.* (2012) Submerged fermentation of the edible mushroom *Pleurotus ostreatus* in a batch stirred tank bioreactor as a promising alternative for the effective production of bioactive metabolites. *Molecules* 17, 2714–2724
- 66 Zhou, X-W. *et al.* (2012) Applied modern biotechnology for cultivation of *Ganoderma* and development of their products. *Appl. Microbiol. Biotechnol.* 93, 941–963
- 67 El Enshasy, H. *et al.* (2010) Bioprocess development for large scale production of anticancer exo-polysaccharide by *Pleurotus ostreatus* in submerged culture. *J. Appl. Sci.* 10, 2523–2529
- 68 Firenzuoli, F. *et al.* (2007) The medicinal mushroom *Agaricus blazei* Murrill: review of literature and pharmaco-toxicological problems. *Evid. Based Complement. Altern. Med.* 5, 3–15
- 69 Geethangili, M. and Tzeng, Y-M. (2011) Review of pharmacological effects of *Antrodia camphorate* and its bioactive compounds. *Evid. Based Complement. Altern. Med.* <http://dx.doi.org/10.1093/ecam/nep108>
- 70 Chen, P.X. *et al.* (2013) Properties of *Cordyceps sinensis*: a review. *J. Funct. Food* 5, 550–569
- 71 Yao, H-Y. *et al.* (2011) *Cytoporus* polysaccharide prevents lipopolysaccharide-induced acute lung injury associated with down-regulating Toll-like receptor 2 expression. *J. Ethnopharmacol.* 137, 1267–1274
- 72 Yin, H. *et al.* (2010) Purification, characterization and immunomodulating properties of polysaccharide isolated from *Flammulina velutipes* mycelium. *Am. J. Chin. Med.* 38, 191–204
- 73 Zhu, X-L. *et al.* (2007) *Ganoderma lucidum* polysaccharides enhance the function of immunological effector cells in immunosuppressed mice. *J. Ethnopharmacol.* 111, 219–226
- 74 Yang, B-K. *et al.* (2007) Chemical characteristics and immunomodulating activities of exo-biopolymers produced by *Grifola frondosa* during submerged fermentation process. *Int. J. Biol. Macromol.* 41, 327–333
- 75 Won, D.P. *et al.* (2011) Immunostimulating activity by polysaccharides isolated from fruiting body of *Inonotus obliquus*. *Mol. Cells* 31, 165–173
- 76 Bisen, P.S. *et al.* (2010) *Lentinus edodes*: a macrofungus with pharmacological activities. *Curr. Med. Chem.* 17, 2419–2430
- 77 Cui, H-L. *et al.* (2011) Isolation, partial characterization and immunomodulatory activities of polysaccharide from *Morchella esculenta*. *J. Sci. Food Agric.* 91, 2180–2185
- 78 Su, C-A. *et al.* (2013) Isolation and characterization of exopolysaccharide with immunomodulatory activity from fermentation broth of *Morchella conica*. *J. Pharm. Sci.* 21, 1–6
- 79 Wu, S-J. *et al.* (2013) *Phellinus linteus* polysaccharides and their immunomodulatory properties in human monocytic cells. *J. Funct. Food* 5, 679–688
- 80 Li, X. and Xu, W. (2011) TLR4-mediated activation of macrophages by the polysaccharide fraction from *Polyporus umbellatus* (*pers.*) *Fries*. *J. Ethnopharmacol.* 135, 1–5
- 81 Cui, F.J. and Chisti, Y. (2003) Polysaccharopeptides of *Coriolus versicolor*: physiological activity, uses and production. *Biotechnol. Adv.* 21, 109–122
- 82 Han, X-Q. *et al.* (2011) Isolation, characterization and immunological activity of a polysaccharide from the fruit bodies of an edible mushroom, *Sarcodon aspratus* (Berk.) S. Ito. *Food Res. Int.* 44, 489–493
- 83 Hobbs, C. (2005) The chemistry, nutritional value, immunopharmacology, and safety of the traditional food of medicinal split-gill fungus *Schizophyllum commune* Fr.:Fr. (Schizophyllaceae). A literature review. *Int. J. Med. Mushrooms* 7, 127–140
- 84 Ohno, N. *et al.* (2002) Antitumor activity and hematopoietic response of a β -glucan extracted from an edible and medicinal mushroom *Sparassis crispa* Wulf:Fr. (Aphyllphoromycetidae). *Int. J. Med. Mushrooms* 4, 13–26
- 85 Ko, H-J. *et al.* (2011) Immunomodulatory properties of *Xylaria nigripes* in peritoneal macrophage cells of Balb/c mice. *J. Ethnopharmacol.* 138, 762–768